

# SEQUENCE LISTING

<110> Taylor, Kent D.  
 Rotter, Jerome I.  
 Yang, Huiying

<120> Methods of Using A Major Histocompatibility Complex  
 Class III Haplotype To Diagnose Crohn's Disease

<130> P-CE 5187

<140>

<141>

<150> US 09/395,345

<151> 1999-09-13

<160> 40

<170> PatentIn Ver. 2.0

<210> 1

<211> 23

<212> DNA

<213> Homo sapiens

<400> 1

ctgtaatccc agctactcaa tcg

23

<210> 2

<211> 24

<212> DNA

<213> Homo sapiens

<400> 2

gggagactta aacagcagaa atgt

24

<210> 3

<211> 21

<212> DNA

<213> Homo sapiens

<400> 3

tcgggatcat ttcagtaatc t

21

<210> 4

<211> 18

<212> DNA

<213> Homo sapiens

<400> 4

gggatccgct gtaactct 18

<210> 5

<211> 22

<212> DNA

<213> Homo sapiens

<400> 5

ggacaatatt ttgctcctga gg 22

<210> 6

<211> 18

<212> DNA

<213> Homo sapiens

<400> 6

gctttgatct cccccctc 18

<210> 7

<211> 24

<212> DNA

<213> Homo sapiens

<400> 7

gagctcggga gtgaggcaga acag 24

<210> 8

<211> 24

<212> DNA

<213> Homo sapiens

<400> 8

tgaggtgtgt tcattagtca actc 24

<210> 9

<211> 21

<212> DNA

<213> Homo sapiens

<400> 9

cgtctctatt tgggcagtga g 21

<210> 10

<211> 18

<212> DNA

<213> Homo sapiens

<400> 10

ggccgaggag gaagaaga

18

<210> 11

<211> 26

<212> DNA

<213> Homo sapiens

<400> 11

ccggatccca taggcctcag agaacc

26

<210> 12

<211> 21

<212> DNA

<213> Homo sapiens

<400> 12

gtaacttaga ttcaggctctg g

21

<210> 13

<211> 20

<212> DNA

<213> Homo sapiens

<400> 13

accaaacttc aaattttcgg

20

<210> 14

<211> 20

<212> DNA

<213> Homo sapiens

<400> 14

gcaacttttc tgtcaatcca

20

<210> 15

<211> 20

<212> DNA

<213> Homo sapiens

<400> 15

ggagctaaaa gttctaactc

20

<210> 16

<211> 20

<212> DNA

Figure 1 consists of 12 line graphs arranged in a 3x4 grid. Each graph plots plasma concentration (mg/ml) on the y-axis against time (hours) on the x-axis. The four groups are labeled (a) through (d) at the top of each column. Each graph contains four data series: control (open circles), diazepam (open squares), diazepam + diazepam (open triangles), and diazepam + diazepam + diazepam (open diamonds). The graphs show that the plasma concentration of diazepam increases over time in all groups, with the highest concentration observed in group (d).

Figure 1 consists of 12 line graphs arranged in a 3x4 grid. Each graph plots plasma concentration (mg/ml) on the y-axis against time (hours) on the x-axis. The four groups are labeled (a) through (d) at the top of each column. Each graph contains four data series: control (open circles), diazepam (open squares), diazepam + diazepam (open triangles), and diazepam + diazepam + diazepam (open diamonds). The graphs show that the plasma concentration of diazepam increases over time in all groups, with the highest concentration observed in group (d).

Figure 1 consists of 12 line graphs arranged in a 3x4 grid. Each graph plots plasma concentration (mg/ml) on the y-axis against time (hours) on the x-axis. The four groups are labeled (a) through (d) at the top of each column. Each graph contains four data series: control (open circles), diazepam (open squares), diazepam + diazepam (open triangles), and diazepam + diazepam + diazepam (open diamonds). The graphs show that the plasma concentration of diazepam increases over time in all groups, with the highest concentration observed in group (d).

Figure 1 consists of 12 line graphs arranged in a 3x4 grid. Each graph plots plasma concentration (mg/ml) on the y-axis against time (hours) on the x-axis. The four groups are labeled (a) through (d) at the top of each column. Each graph contains four data series: control (open circles), diazepam (open squares), diazepam + diazepam (open triangles), and diazepam + diazepam + diazepam (open diamonds). The graphs show that the plasma concentration of diazepam increases over time in all groups, with the highest concentration observed in group (d).

Figure 1 consists of 12 line graphs arranged in a 3x4 grid. Each graph plots plasma concentration (mg/ml) on the y-axis against time (hours) on the x-axis. The four groups are labeled (a) through (d) at the top of each column. Each graph contains four data series: control (open circles), diazepam (filled circles), diazepam + diazepam (open squares), and diazepam + diazepam + diazepam (filled triangles). The graphs show that the plasma concentration of diazepam increases over time in all groups, with the highest concentration observed in group (d).

Figure 1 consists of 12 line graphs arranged in a 3x4 grid. Each graph plots 'Plasma concentration (mg/ml)' on the y-axis (0 to 12) against 'Time (h)' on the x-axis (0 to 12). The graphs are labeled (a) through (l). The first row (a, b, c, d) shows the effect of 100 mg/kg of diazepam in control, diazepam, diazepam + diazepam, and diazepam + diazepam + diazepam groups, respectively. The second row (e, f, g, h) shows the effect of 100 mg/kg of diazepam in control, diazepam, diazepam + diazepam, and diazepam + diazepam + diazepam groups, respectively. The third row (i, j, k, l) shows the effect of 100 mg/kg of diazepam in control, diazepam, diazepam + diazepam, and diazepam + diazepam + diazepam groups, respectively. The graphs show that the plasma concentration of diazepam increases over time, with the highest concentrations observed in the diazepam + diazepam + diazepam group (graphs d, h, l).

Figure 1 consists of 12 line graphs arranged in a 3x4 grid. Each graph plots 'Plasma concentration of diazepam (mg/ml)' on the y-axis (ranging from 0 to 12) against 'Time (min)' on the x-axis (ranging from 0 to 120). The four groups are labeled (a) through (d) at the top of each column. Each graph shows a single data series with a peak around 60 minutes. The peak heights are approximately: (a) 4.5, (b) 6.5, (c) 8.5, and (d) 10.5 mg/ml. The curves are smooth and bell-shaped, indicating a rapid increase followed by a gradual decline.

Figure 1 consists of 12 line graphs arranged in a 3x4 grid. Each graph plots plasma concentration (mg/ml) on the y-axis (0 to 12) against time (hours) on the x-axis (0 to 12). The four groups are: (a) control, (b) diazepam, (c) diazepam + diazepam, and (d) diazepam + diazepam. The graphs show that the plasma concentration of diazepam increases over time in all groups, with the highest concentration observed in group (d) at 12 hours.

<213> Homo sapiens

<400> 22

tgagacagag gataggagag acag

24

<210> 23

<211> 24

<212> DNA

<213> Homo sapiens

<400> 23

ggtttctctg actgcatctt gtcc

24

<210> 24

<211> 23

<212> DNA

<213> Homo sapiens

<400> 24

tcatggggag aacctgcaga gaa

23

<210> 25

<211> 22

<212> DNA

<213> Homo sapiens

<400> 25

cctctctccc ctgcaacaca ca

22

<210> 26

<211> 24

<212> DNA

<213> Homo sapiens

<400> 26

gcctctagat ttcattccagc caca

24

<210> 27

<211> 22

<212> DNA

<213> Homo sapiens

<400> 27

gtgtgtgttg caggggagag ag

22

<210> 28

<211> 22

<212> DNA

<213> Homo sapiens

<400> 28

gcactccagc ctaggccaca ga

22

<210> 29

<211> 21

<212> DNA

<213> Homo sapiens

<400> 29

acaatggaca cttgggttac t

21

<210> 30

<211> 19

<212> DNA

<213> Homo sapiens

<400> 30

aatgagatgc cacctgaaa

19

<210> 31

<211> 24

<212> DNA

<213> Homo sapiens

<400> 31

ttatctactt atagtctatc acgg

24

<210> 32

<211> 24

<212> DNA

<213> Homo sapiens

<400> 32

ggcttgactt gaaactcaga gacc

24

<210> 33

<211> 25

<212> DNA

<213> Homo sapiens

<400> 33

gatcgctcac cagcacactg gctat

25

<210> 34

<211> 25

<212> DNA

<213> Homo sapiens

<400> 34

ctgggcaaca gagcgagctc cgtct

25

<210> 35

<211> 22

<212> DNA

<213> Homo sapiens

<400> 35

tccttacagc agagatatgt gg

22

<210> 36

<211> 22

<212> DNA

<213> Homo sapiens

<400> 36

agatggcatt tggagagtgc ag

22

<210> 37

<211> 21

<212> DNA

<213> Homo sapiens

<400> 37

caacacactg atttccatag c

21

<210> 38

<211> 21

<212> DNA

<213> Homo sapiens

<400> 38

aatgggctac tacttcacac c

21

<210> 39

<211> 21

<212> DNA

<213> Homo sapiens

<400> 39

gaaatgtgag aataaaggag a

21

<210> 40

<211> 20

<212> DNA

<213> Homo sapiens

<400> 40

gataaagggg aactactaca

20

gataaagggg aactactaca